

# Cross-Sectional Comparison of Executive Attention Function in Normally Aging Long-Term *T'ai Chi*, Meditation, and Aerobic Fitness Practitioners Versus Sedentary Adults

Teresa D. Hawkes, PhD,<sup>1</sup> Wayne Manselle, MS,<sup>2</sup> and Marjorie H. Woollacott, PhD<sup>2</sup>

## Abstract

This cross-sectional field study documented the effect of long-term *t'ai chi*, meditation, or aerobic exercise training versus a sedentary lifestyle on executive function. It was predicted that long-term training in *t'ai chi* and meditation plus exercise would produce greater benefits to executive function than aerobic exercise. *T'ai chi* and meditation plus exercise include mental and physical training. Fifty-four volunteers were tested: *t'ai chi* ( $n=10$ ); meditation+exercise ( $n=16$ ); aerobic exercisers ( $n=16$ ); and sedentary controls ( $n=12$ ). A one-factor (group), one-covariate (age) multivariate analysis of covariance was performed. Significant main effects of group and age were found (group, 67.9%,  $p<0.001$ ; age, 76.3%,  $p=0.001$ ). *T'ai chi* and meditation practitioners but not aerobic exercisers outperformed sedentary controls on percent switch costs ( $p=0.001$  and  $p=0.006$ , respectively), suggesting that there may be differential effects of training type on executive function.

## Introduction

AS HUMAN LIFE EXPECTANCY has lengthened in the developed world, successful aging has become a public health concern. Quality of life, cognitive capacity, and physiologic status are all affected by aging.<sup>1</sup> Aging effects include declines in key physiologic systems (cardiovascular, neuromotor, and cognitive), and complicating effects of damage to those systems. Maximal oxygen uptake (VO<sub>2</sub> Max), a well-established proxy for cardiovascular health, declines at a rate of 1% per year after the age of 20.<sup>2</sup> Reaction time (RT) on neuropsychologic tests of cognitive capacity has been shown to decline by a factor of 1.5 during the 25th–65th years of life.<sup>3</sup> Furthermore, evidence suggests that the seventh decade is a critical point when significantly greater decrements begin to occur.<sup>4,5</sup> After the age of 65, walking and other locomotor skills necessary for independent, daily living require increasing cognitive resources to perform.<sup>6</sup> Can these declines be mitigated?

This cross-sectional field study observed the aerobic and executive function capacity of self-selected long-term exercisers in three groups (*t'ai chi*, meditation plus exercise, and aerobic exercise) versus sedentary controls living in Eugene, Oregon. Investigation of training regimens that may offset the effects of aging on physiologic function has been under way for decades. Of particular interest are regimens that may

extend mid-life (30–65 years) cognitive capacity into the seventh and eighth decades of life. Aerobic exercise and meditation are two available health regimens shown to positively affect executive function in young and older adults.<sup>7–13</sup>

Executive function processes involved in tasks of daily living include the ability to respond appropriately to novel situations, make choices, set goals, coordinate and modify task sequences, inhibit inappropriate responses, and switch between tasks or instructions appropriately.<sup>4,14</sup> Key executive components underlying these operations are (1) inhibition, (2) updating, (3) shifting between instruction sets,<sup>15</sup> and working memory.<sup>14</sup> Salthouse et al.<sup>4</sup> found that degradation of the inhibition and updating components of executive function may mediate age-related cognitive decline (i.e., memory, percept formation, association processes).

A small number of studies have divided exercise types into those requiring aerobic exertion alone and aerobic exertion plus consistent mental exertion. Exercise requiring consistent mental exertion (e.g., soccer, dance) was associated with better performance on tests of executive attention<sup>16,17</sup> Exercise requiring consistent mental exertion generally also requires cortical coordination of complex movement sequences.<sup>18</sup>

Aspects of executive function that are upregulated by moderate exercise include speed of information processing

<sup>1</sup>University of Oklahoma, Norman, OK.

<sup>2</sup>University of Oregon, Eugene, OR.

as indexed by visual and auditory electroencephalographic (EEG) P300 event-related potential (ERP) latency and reaction-time decreases, and correlated accuracy increases in oddball tasks; and (2) enhancement of executive attention processes, indexed by RT and accuracy on Erikson flanker tasks in older adults.<sup>7</sup> Additionally, research shows that consistent aerobic exercise regimens increase the thickness of frontal, parietal, and temporal cortices in humans.<sup>19</sup> A recent meta-analysis showed that intense aerobic exercise as quantified by standard cardiovascular measures is not required to accrue cognitive benefits from exercise. Light-to-moderate exercise and dose frequency are key factors.<sup>20</sup>

Meditation has also been shown to enhance executive function. Meditation is defined as concentrated mental focus on a sound, image, set of syllables (chant or mantra), or activity (walking, sitting). All conflicting stimuli are pushed out of awareness.<sup>21</sup> Thus, meditation requires mental effort to inhibit distracting thoughts or sensations. Chan and Woollacott<sup>10</sup> found that long-term practitioners of meditation showed less interference on the Stroop task than nonpracticing controls. Importantly, many meditation practitioners also engage in yoga and other moderate exercise activities. Indeed, one of the most utilized and studied meditation practices, mindfulness-based stress reduction, explicitly incorporates Hatha yoga into its training regimen. Hatha yoga is a form of moderate exercise requiring complex coordination of body segments.<sup>22</sup> Not surprisingly, during subject recruitment for this study, no sedentary meditators were found. All meditation practitioners reported habitual participation in moderate exercise of some kind. Thus, a contributing factor in meditation's benefit to executive function might be cardiovascular and metabolic modifications resulting from a lifestyle typically including moderate aerobic exertion.

Another health training regimen that shows promise for benefiting executive function is *t'ai chi*. *T'ai chi* is a form of moderate exercise<sup>23</sup> that has been shown to be superior to aerobic walking for cardiovascular function.<sup>24</sup> *T'ai chi* is also a form of moving meditation<sup>25</sup> requiring memorization of complex movement sequences,<sup>26</sup> and thus motor learning. Motor learning requires executive function.<sup>27</sup> *T'ai chi* also requires constant application of attention for optimal performance.<sup>26</sup> Because *t'ai chi* requires moderate exercise, motor learning, consistent application of attention, and complex coordination of movement, it is possible that frequent practice would yield similar or greater benefits to executive function compared to aerobic exercise alone. Indeed, a recent uncontrolled study showed that 10 weeks of *t'ai chi* training was associated with better task-switch performance in community-dwelling elders.<sup>28</sup>

This cross-sectional field study of long-term *t'ai chi*, meditation, and aerobic exercise practitioners versus sedentary controls utilized a complex visuospatial task switch (VSTS) test to assess executive function. This test required three main components of executive function: (1) inhibition, (2) updating, and (3) shifting between instruction sets. Subjects were also tested with the Rockport 1-mile walk, a well-validated assessment of estimated VO<sub>2</sub> Max, to document cardiovascular status.<sup>29</sup> It was expected that the training groups would outperform sedentary controls on estimated VO<sub>2</sub> Max. As a result, it was also expected they would outperform sedentary controls on executive function measures.<sup>7,30,31</sup> It was predicted that long-term *t'ai chi* and

meditation practice would be associated with greater benefits to executive function than aerobic exercise alone, because both combine mental and physical exertion. All three training groups were predicted to significantly outperform sedentary controls on executive function. Further, it was hypothesized that age would have less effect on this study's key measures in health regimen practitioners compared to sedentary controls.

## Materials and Methods

### Participants

Participants were recruited from Eugene, Oregon. Inclusion criteria were (1) no self-reported neurological or physical disorders, (2) living independently, and (3) aged 20–75. The large age range was chosen so the effects of aging could be documented over the adult lifespan on the key measures. It was expected that group membership would be associated with larger effects than age.<sup>3,5,8,32–34</sup> Sedentary participants were required to have (1) a generally inactive lifestyle for 5 or more years, and (2) no prior experience with meditation or *t'ai chi*. Training group practitioners were required to (1) have practiced at least 5 years or more, three times per week, 30 minutes per session. All participants had self-selected into their preferred level and type of exercise activity. Fifty-nine participants enrolled in the study. Because acute exercise has been shown to improve cognitive performance,<sup>17,35</sup> cognitive testing was done before exercise testing. Two participants unable to use a computer effectively were excluded, since the executive attention test was administered via computer. Two subjects did not complete the testing. One subject who presented with bipolar disorder was excluded. Fifty-four subjects were thus included in this analysis (female=27). Group composition was (1) 10 *t'ai chi* (female=3), (2) 16 meditation plus exercise (female=6), (3) 16 aerobic fitness (female=8), and (4) 12 sedentary (female=10) participants. Body-mass index (BMI), which has been associated with cognitive capacity,<sup>36,37</sup> was calculated for each participant. BMI is mass in kilograms/height in meters squared (Table 1).<sup>2</sup> Participant recruitment and experimental protocol were approved by the University of Oregon Institutional Review Board. Participants gave informed consent and were compensated for participation.

### Testing

**Aerobic capacity:** Rockport 1-mile walk.<sup>29</sup> The Rockport 1-mile walk was utilized to obtain estimated VO<sub>2</sub> Max, controlling for age, weight, and gender (<http://www.exrx.net/Calculators/Rockport.html>).<sup>29</sup>

**Executive attention.** Executive attention test structure has been shown to affect RT. To maximize test difficulty, a complex visuospatial task switch test (Mayr Laboratory, University of Oregon, 2009) was utilized. This test required working memory, inhibition, shifting, and updating,<sup>15</sup> was noncued, and utilized alternating runs of two rules.<sup>38</sup> Stimulus appearance was randomized. More complex tests produce longer RTs.<sup>32</sup> Noncued paradigms have been shown to be more difficult than cued paradigms.<sup>39</sup> Short response-stimulus intervals have been shown to produce greater switch costs than longer response to stimulus intervals.<sup>40</sup> The

TABLE 1. PARTICIPANT PHYSIOLOGIC SCORES

Group	n	Females	Age		VO <sub>2</sub> Max		BMI	
			M	SD	M	SD	M	SD
T'ai chi	10	3	55.4	12.99	34.14	6.34	29.3	3.77
Meditation	16	6	48.63	15.00	41.83	9.04	23.3	3.53
Aerobics	16	8	44.09	16.2	45.66	9.67	23.78	2.62
Sedentary	12	2	46.92	12.81	28.68	5.76	27.93	6.37

Global ranges: age, 22–75; BMI, 18.50–37.90; estimated VO<sub>2</sub> Max, 17.23–60.00.

VO<sub>2</sub> Max, estimated maximal oxygen uptake; BMI, body-mass index; SD, standard deviation; M, mean.

Mayr VSTS utilized a short response to stimulus interval (10msec). Measures were switch RT and switch capacity (percent local switch costs). Percent local switch costs are calculated by subtracting no-switch from switch RT, then dividing by no-switch RT to control for the speed accuracy trade-off. Local switch costs index the capacity to switch tasks quickly and accurately.<sup>39,41,42</sup> Lower switch costs index greater switch capacity.<sup>17</sup>

**Paradigm.** Participants were trained to indicate the location of a randomly appearing dot within a fixation rectangle using two different response rules (Fig. 1). For Rule 1, the button-press response was compatible with the dot's location in space, but for Rule 2, was incompatible. For Switch trials (Rule 3), participants switched between Rule 1 and 2 on every other trial.

Stimuli were displayed on a computer monitor located ~24 inches in front of the participant. Participants were trained to respond as quickly and accurately as possible. For Rule 3, participants were provided with visual feedback regarding errors. They corrected their error and continued the trial block. Participants practiced until they achieved 85% accuracy. Rules 1 and 2 consisted of 48 trials in two blocks. Rule 3 (the actual task switch) consisted of 12 blocks of 48 trials/block.

**Multivariate cross-sectional design.** In multivariate designs, multiple dependent variables are measured on subjects




FIG. 1. Visuospatial task switch paradigm (Mayr Laboratory, University of Oregon).

who are assigned membership in carefully defined groups.<sup>43</sup> The study was a one-factor (group), one-covariate (age) multivariate analysis of covariance (MANCOVA) with groups: (1) *t'ai chi*, (2) meditation+exercise, (3) aerobic exercise, and (4) sedentary controls who had never engaged in the training modalities. Age and group were expected to have significant effect sizes on VO<sub>2</sub> Max, switch reaction time, and percent local switch costs. Each training group reported statistically similar lifetime hours of moderate aerobic exertion. This allowed the investigators to equate physical exercise effects. Covarying age allowed assessing its effect on the dependent measures. The difference between the training groups was type of attentional focus and complexity of motor output required.<sup>16,17</sup>

### Data analysis

A multivariate analysis of covariance (MANCOVA) with one factor (group) and one covariate (age), and Levene's test for homogeneity of variance were performed. Alpha was 0.05 for the MANCOVA. A bivariate correlation was run on all variables. To control for  $\alpha$  slippage for multiple analyses, a Bonferroni correction was applied and  $\alpha$  was set at 0.0125. Dependent measures were (1) estimated VO<sub>2</sub> Max, (2) switch reaction time, and (3) percent local switch costs. Because whole system functionality was evaluated, error trials were included in reaction time means. To document error effects on RT, mean RTs of error trials (milliseconds), accurate trials (milliseconds), and post-error trials (milliseconds) were compared in a one-way analysis of variance (ANOVA) with trial type as factor and trial time in milliseconds as the dependent measure. To isolate predictors of switch costs, a regression with group, age, BMI, VO<sub>2</sub> Max, error trials, accurate trials, percent accuracy, and post-error trials as predictors of percent local switch costs was run. It was predicted that all training groups would outperform sedentary controls. All analyses were run with PSAW Statistics 19 (IBM, Chicago, IL).

### Results

#### Overall results

As expected, training group and age showed significant effect sizes on key outcome measures. The main MANCOVA omnibus was significant (Wilk's lambda ( $\Lambda$ ) ( $F(24, 26)=1417.561, p<0.001$ .) Our overall partial  $\eta^2$  was .999, indicating that 99.9% of the variance in outcome measures were explained. Since key variables related to executive function—aerobic capacity and age—were included, this is not surprising. Group membership explained ~68% of variance (Wilk's lambda ( $\Lambda$ ) ( $F(72, 78.562)=2.321, p<0.001$ , partial  $\eta^2=.679$ ), indicating the presence of possible training effects. Though groups did not differ on age ( $p=0.295$ ), age explained 76% of total variance (Wilk's lambda ( $\Lambda$ )  $F(24, 26)=3.488, p=0.001$ , partial  $\eta^2=.763$ ), suggesting that normal aging was a key factor affecting the outcome. Clearly these two have overlapping variance, as would be expected, since the cell populations that produce the output are affected by both age and training effects.<sup>7,30,33</sup> Means, standard deviations, and variance explained for key factors and variables are presented in Figure 2. Age and group membership had similar effect sizes on switch RT (group, 34.3%; age, 31.2%) and VO<sub>2</sub> Max (group,




FIG. 2. Effect-size profiles. Effect of group and age on key system variables. RT, reaction time; VO<sub>2</sub> Max, maximal oxygen uptake; BMI, body-mass index.

49.6%; age, 39.3%), but not on percent switch costs (group, 28%; age, 9%) (Fig. 2, Table 2, and Table 3). Error trials ( $\beta = .012$ ,  $t(44) = -.520$ ,  $p > 0.05$ ) did not significantly predict percent local switch costs and were included in the analysis. Mean accuracy was not significantly different between the groups ( $p = 0.425$ ). A one-way ANOVA with BMI as the dependent variable and group as the factor showed significant differences between groups ( $F(3, 50) = 6.569$ ,  $p = 0.001$ ).

**Post-hoc results.** Our groups did not differ on age (Fig. 3a). *T'ai chi* ( $p = 0.025$ ), meditation ( $p < 0.001$ ), and aerobic fitness ( $p < 0.001$ ) practitioners outperformed sedentary controls on estimated VO<sub>2</sub> Max. Aerobic exercisers outperformed *t'ai chi* practitioners ( $p = 0.043$ ) on VO<sub>2</sub> Max (Fig. 3a). This may be due to the wide range of exertion required by different *t'ai chi* styles (i.e., Chen versus Yang style, long-versus short-form).<sup>23,44</sup> VO<sub>2</sub> Max was negatively correlated with age ( $r = -.539$ ,  $p < 0.001$ ). BMI differed between groups ( $p = 0.002$ ). *T'ai chi* and sedentary controls showed significantly higher BMI than meditators ( $p = 0.004$  and  $p = 0.036$ , respectively). *T'ai chi* BMI was significantly higher than

TABLE 2. PARTICIPANT COGNITIVE SCORES

Group	n	SwRT		SwCosts	
		M	SD	M	SD
<i>T'ai chi</i>	10	453.94	110.84	14.13	11.97
Meditation	16	477.41	188.88	401.23	122.83
Aerobics	16	489.63	96.89	400.84	53.90
Sedentary	12	654.3	154.56	654.3	154.56

Global ranges: switch reaction time, 301.88–1104.1; % local switch costs, –0.19 to 57.87.

SwRT, switch reaction time (msec); SwCosts, % local switch costs.

TABLE 3. EFFECT SIZE BY GROUP, AGE, AND SUBJECT MEASURES

IV, CoV	Physiologic measures		Cognitive measures	
	Est. VO <sub>2</sub> Max	BMI	SwRT	SwCosts
IV-Group	49.60**	29.00*	34.30**	28.20*
CoV-Age	39.30*	1.10	31.20**	9.30*

\* $p < 0.05$ . \*\* $p < 0.001$ .

Effect size = partial  $\eta^2$ .

IV, quasi-independent variable; CoV, covariate; Est. VO<sub>2</sub>, estimated VO<sub>2</sub> Max (mL O<sub>2</sub>/kg/min); BMI, body-mass index (kg/m<sup>2</sup>); SwRT, switch reaction trial time (RT) (msec); SwCosts, % local switch costs (switch RT – no-switch RT/no-switch RT).

aerobic exercisers ( $p = 0.009$ ). Aerobic exercisers and sedentary controls did not differ on BMI.

*T'ai chi* and meditation groups outperformed sedentary controls on percent switch costs: *t'ai chi*:  $p = 0.001$ ; meditators:  $p = 0.006$ . Aerobic exercisers and sedentary controls did not differ (Fig. 3c). Training groups outperformed sedentary




FIG. 3. Key observations by group. (a) Age covariate. (b) Estimated maximal oxygen uptake (VO<sub>2</sub> Max). (c) Body-mass index (BMI). (d) Visuospatial task switch (VSTS) costs. (e) VSTS reaction time. Error bars are  $\pm 1$  standard error (SE). TC, *t'ai chi*; M, meditation; A, aerobic; S, sedentary.

TABLE 4. CORRELATIONS BETWEEN KEY MEASURES

Grp	Age	VO <sub>2</sub>	BMI	SwRT	SwCosts
Grp	-.206	-.142	-.037	.400*	.468*
Age		-.539**	.003	.433*	.181
VO <sub>2</sub>			-.386*	-.508**	-.289
BMI				.144	-.010
SwRT					.660**
SwCosts					

\* $p < 0.0125$ , \*\* $p < 0.001$ .

Grp, group (*t'ai chi*=1, meditation=2, aerobic=3, sedentary=4); Age (years); estimated VO<sub>2</sub> Max, estimated maximal oxygen uptake (mL O<sub>2</sub>/kg/min); BMI, body-mass index (kg/m<sup>2</sup>); SwRT, switch reaction trial time (msec); SwCosts, % local switch costs (switch RT-no-switch RT/no-switch RT).

controls on switch RT: *t'ai chi* ( $p < 0.001$ ), meditation ( $p = 0.001$ ), aerobic exercisers ( $p = 0.014$ ) (Fig. 3d). There were no differences between training groups on switch RT. Group was significantly and positively correlated with both switch RT ( $p = 0.003$ ,  $r = .400$ ) and percent switch costs ( $p < 0.001$ ;  $r = .468$ ) (Table 4). Thus, aerobic exercisers and sedentary controls demonstrated longer RTs and higher switch costs than *t'ai chi* and meditation practitioners. Figure 4 shows z-score comparisons by key variable.

## Discussion

This cross-sectional field study of the effects of long-term *t'ai chi*, meditation+exercise, or aerobic exercise training versus a sedentary lifestyle utilized a VSTS test<sup>5</sup> to assess complex executive function. It was hypothesized that self-

reported long-term practice of these modalities would be inversely associated with local switch costs: an index of executive function. First it was asked if the groups differed on aerobic capacity. If they did, those with higher VO<sub>2</sub> Max could be expected to outperform individuals with lower VO<sub>2</sub> Max on executive and cardiovascular indices.

It was then asked if groups differed on key executive function measures: switch RT and switch capacity (percent local switch costs). It was predicted that (1) training groups would outperform sedentary controls on estimated VO<sub>2</sub> Max, and (2) long-term *t'ai chi* and meditation plus exercise training would produce the greatest benefits to the executive function measures. These would be followed by aerobic exercisers, then sedentary controls. Furthermore, we hypothesized that age effects would be offset by training effects.

As expected, each of our training groups outperformed sedentary controls on estimated VO<sub>2</sub> Max. RT and switch cost benefits conferred by long-term moderate exercise were associated with the training groups, but not with sedentary controls. Even though groups were equated on age, and age significantly affected all outcome measures, post-hoc results showed that these effects fell most detrimentally on the sedentary controls: slower RT, higher switch costs, and lower VO<sub>2</sub> Max. Furthermore, age should have affected executive function equally in each group,<sup>7</sup> yet the training groups outperformed sedentary controls when controlling for age. This suggests that the executive function differences between groups may be associated with training effects. In line with previous studies on switch RT,<sup>7</sup> all the training groups outperformed sedentary controls. However, on the more stringent switch costs measure, only *t'ai chi* and meditation groups significantly outperformed sedentary controls.

As noted, even though groups did not differ significantly on age ( $p = 0.295$ ), age was associated with a large effect size in outcome measures. Age effects generally went in the direction expected: longer RTs and lower cardiovascular capacity (estimated VO<sub>2</sub> Max). However, an inspection of the distribution of age scores by group showed *t'ai chi* and meditation groups were older than the aerobic fitness or sedentary groups, so larger age effects associated with these two groups' scores should be seen. However, on average, younger aerobic fitness practitioners demonstrated higher switch costs than older *t'ai chi* and meditation practitioners. Finally, group was positively correlated with switch costs ( $p < .001$ ,  $r = .468$ ). *T'ai chi* was dummy coded 1, meditation 2, aerobic fitness 3, and sedentary controls, 4. These results show that higher group number is correlated with higher switch costs. This convergent evidence suggests that moderate aerobic exercise requiring consistent mental exertion is associated with greater benefits to attentional tasks requiring coordination of inhibition, updating, shifting, and working memory than aerobic exercise alone.

However, limitations in the data set could contribute to this outcome. The sample size of this study is small. Therefore, a lack of statistical power could contribute to the modest differences between the aerobic and sedentary groups on switch RT. There was no specific screening for mild cognitive disorders, which may have impacted the neuropsychologic outcomes. Self-selection into these groups related to socio-economic status, and alleles of moderators of executive function such as catechol-O-methyltransferase (COMT) and brain-derived neurotrophic factor (BDNF),<sup>30,31,45</sup> could be driving the results.




FIG. 4. Possible training effects on physiologic and cognitive capacity by key variables. Error bars are  $\pm 1$  standard error (SE). VO<sub>2</sub> Max, estimated maximal oxygen uptake; RT, reaction time; T, *t'ai chi*; M, meditation; A, aerobic; S, sedentary.

Finally, though data were collected on which type of exercise or *t'ai chi* each person practiced, the samples collected of each were not large enough to generalize these outcomes to the larger population of each of these types. Certainly different intensities of practice across participants could affect the outcome measures. Indeed, the literature suggests that VO<sub>2</sub> Max is differentially affected by exercise intensity (i.e., low-to-high-exertion requirements).<sup>46</sup> A longitudinal study comparing training effects on VO<sub>2</sub> Max and executive attention correlated with BDNF and COMT alleles could begin to address some of these limitations.

In spite of these limitations, this evidence suggests that practice of *t'ai chi* and meditation+exercise may be associated with equal or better performance on a stringent measure of executive function than practice of aerobic fitness alone. This is good news for clinicians and individuals alike. Health care professionals need different types of evidence-based health practices to offer clients. If clients are able to select from regimens based on personal inclinations and preferences, it is possible that participation in such regimens would become habitual, ensuring cognitive and physical benefits, lower health care costs, and greater quality of life into later years.

### Acknowledgments

This work was supported by a Francisco J. Varela Research Award, Mind & Life Institute, 2007, and an NIH T-32 Systems Training Grant Appointment, Institute of Neuroscience, University of Oregon, 2008. Andrew Lovering, PhD contributed valuable advice on estimated VO<sub>2</sub> Max.

### Disclosure Statement

No competing financial interests exist.

### References

1. Murdaugh C. Health-related quality of life as an outcome in organizational research. *Med Care* 1997;35:N541–N548.
2. Powers SK, Howley ET. *Exercise Physiology: Theory and Application to Fitness and Performance*, 5<sup>th</sup> edition. Boston: McGraw Hill, 2004.
3. Verhaeghen P, Cerella J. Aging, executive control, and attention: A review of meta-analyses. *Neurosci Biobehav Rev* 2002;26:849–857.
4. Salthouse TA, Atkinson TM, Berish DE. Executive functioning as a potential mediator of age-related cognitive decline in normal adults. *J Exp Psychol Gen* 2003;132:566–594.
5. Mayr U. Age differences in the selection of mental sets: The role of inhibition, stimulus ambiguity, and response-set overlap. *Psychol Aging* 2001;16:96–109.
6. Shumway-Cook A, Woollacott M. *Motor Control: Translating Research into Clinical Practice*. Philadelphia: Lippincott, Williams & Wilkins, 2011.
7. Ratey JJ, Loehr JE. The positive impact of activity on cognition during adulthood: A review of underlying mechanisms, evidence and recommendations. *Rev Neurosci* 2011;22:171–195.
8. Etnier JL, Chang YK. The effect of physical activity on executive function: A brief commentary on definitions, measurement issues, and the current state of the literature. *J Sport Exerc Psychol* 2009;31:469–483.
9. Tang SW, Chu E, Hui T, et al. Influence of exercise on serum brain-derived neurotrophic factor concentrations in healthy human subjects. *Neurosci Lett* 2008;431:62–65.
10. Chan D, Woollacott MH. Effects of level of meditation experience on attentional focus: Is the efficiency of executive or orientation networks improved? *J Altern Complement Med* 2007;36:651–657.
11. Brefczynski-Lewis JA, Lutz A, Schaefer HS, et al. Neural correlates of attentional expertise in long-term meditation practitioners. *Proc Natl Acad Sci* 2007;104:11483–11488.
12. Slagter HA, Lutz A, Greischar LL, et al. Mental training affects distribution of limited brain resources. *PLoS Biol* 2007;5:1228–1235.
13. Hölzel BK, Carmody J, Vangel M, et al. Mindfulness practice leads to increases in regional brain gray matter density. *Psychiatry Res* 2011;191:36–43.
14. Gilbert SJ, Burgess PW. Executive function. *Curr Biol* 2008;18:R110–R114.
15. Miyake A, Friedman NP, Emerson MJ, et al. The unity and diversity of executive functions and their contributions to complex “frontal lobe” tasks: A latent variable analysis. *Cogn Psychol* 2000;41:49–100.
16. Voss MW, Kramer AF, Basak C, et al. Are expert athletes ‘expert’ in the cognitive laboratory? A meta-analytic review of cognition and sport exercise. *Appl Cogn Psychol* 2009;24:812–826.
17. Pesce C, Audiffren M. Does acute exercise switch off switch costs? A study with younger and older athletes. *J Sport Exerc Physiol* 2011;33:609–626.
18. Blasing B, Puttke M, Schack, T. *The Neurocognition of Dance: Mind, Movement and Motor Skills*. New York: Psychology Press, 2010.
19. Kramer AF, Colcombe SJ, McAuley E, et al. Fitness, aging and neurocognitive function. *Neurobiol Aging* 2005;26S:S124–S127.
20. Etnier JL, Nowell PM, Landers DM, Sibley BA. A meta-regression to examine the relationship between aerobic fitness and cognitive performance. *Brain Res Rev* 2006;52:119–130.
21. Lutz A, Dunne JD, Davidson RJ. Meditation and the neuroscience of consciousness: An introduction. In: Zelazo PD, Moscovitch M, Thompson E, eds. *The Cambridge Handbook of Consciousness*. Cambridge, UK: Cambridge University Press, 2007.
22. Smith BW, Shelley BM, Dalen J, et al. A pilot study comparing the effects of mindfulness-based and cognitive-behavioral stress reduction. *J Altern Complement Med* 2008;14:251–258.
23. Li JX, Hong Y, Chan KM. Tai chi: Physiological characteristics and beneficial effects on health. *Br J Sports Med* 2001;35:148–156.
24. Audette JF, Jin YS, Newcomer R, et al. Tai chi versus brisk walking in elderly women. *Age Aging* 2006;35:388–393.
25. Luskin F. Transformative practices for integrating mind-body-spirit. *J Altern Complement Med* 2004;10(Suppl 1):S15–S23.
26. Gatts S. A Tai Chi Chuan training model to improve balance control in older adults. *Curr Aging Sci* 2008;1:68–70.
27. Halsband U, Lange RK. Motor learning in man: A review of functional and clinical studies. *J Physiol Paris* 2006;99:414–424.
28. Matthews MM, Williams HG. Can Tai chi enhance cognitive vitality? A preliminary study of cognitive executive control in older adults after a Tai chi intervention. *J S C Med Assoc* 2008;104:255–257.
29. Kline GM, Porcari JP, Hintermeister R, et al. Estimation of VO<sub>2</sub> Max from a one mile track walk, gender, age, and body weight. *Med Sci Sports Exerc* 1987;19:253–259.

30. Vaynman S, Gomez-Pinilla F. Revenge of the "sit": How lifestyle impacts neuronal and cognitive health through molecular systems that interface energy metabolism with neuronal plasticity. *J Neurosci Res* 2006;84:699–716.
31. Dishman RK, Berthoud HR, Booth FW, et al. The neurobiology of exercise. *Obesity* 2006;14:345–356.
32. Bryan J, Luszcz MA. Measurement of executive function: Considerations for detecting adult age differences. *J Clin Exp Neuropsychol* 2000;22:40–55.
33. Hillman HH, Weiss EP, Hagberg JM, Hatfield BD. The relationship of age and cardiovascular fitness to cognitive and motor processes. *Psychophysiology* 2002;39:303–312.
34. Colcombe SJ, Kramer AF, Erickson KI, et al. Cardiovascular fitness, cortical plasticity, and aging. *Proc Natl Acad Sci* 2004;101:3316–3321.
35. Davranche K, Audiffren M. Facilitating effects of exercise on information processing. *J Sports Sci* 2004;22:419–428.
36. Gunstad J, Lhotsky A, Wendell CR, et al. Longitudinal examination of obesity and cognitive function: Results from the Baltimore longitudinal study of aging. *Neuroepidemiology* 2010;34:222–229.
37. Sabia S, Nabi H, Kivimaki M, et al. Health behaviors from early to late midlife as predictors of cognitive function: The Whitehall II study. *Am J Epidemiol* 2009;170:428–437.
38. Altmann EM, Gray WD. An integrated model of cognitive control in task switching. *Psychol Rev* 2008;115:602–639.
39. Monsell S, Sumner P, Waters H. Task-set reconfiguration with predictable and unpredictable task switches. *Mem Cognit* 2003;31:327–342.
40. Karayanidis F, Coltheart M, Michie PT, Murphy K. Electrophysiological correlates of anticipatory and poststimulus components of task-switching. *Psychophysiology* 2003;40:329–348.
41. Wylie G, Allport A. Task switching and the measurement of "switch costs." *Psychol Res* 2000;63:212–233.
42. Milan EG, Gonzalez A, Sanabria D, et al. The nature of residual cost in regular switch response factors. *Acta Psychol (Amst)* 2006;122:45–57.
43. Stevens JP. *Applied Multivariate Statistics for the Social Sciences*, 4<sup>th</sup> edition. Mahwah, NJ: Lawrence Erlbaum Associates, 2002.
44. Wolf SL, Coogler C, Tingsen X. Exploring the basis for Tai Chi Chuan as a therapeutic exercise approach. *Arch Phys Med Rehabil* 1997;78:886–892.
45. Gajewski PD, Hengstler JG, Golka K, et al. The met-allele of the BDNF Val66Met polymorphism enhances task switching in elderly. *Neurobiol Aging* 2011;32:2327.
46. Tanaka H, DeSouza CA, Jones PP, et al. Greater rate of decline in maximal aerobic capacity with age in physically active vs. sedentary healthy women. *J Appl Physiol* 1997;83:1947–1953.

Address correspondence to:  
Teresa D. Hawkes, PhD  
3730 West Amazon Drive  
Eugene, OR 97405

E-mail: [teresa.hawkes@gmail.com](mailto:teresa.hawkes@gmail.com)

Copyright of Journal of Alternative & Complementary Medicine is the property of Mary Ann Liebert, Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.